

**CLAIMS**

1. A polypeptide which:
  - (i) comprises the amino acid sequence as recited in SEQ ID NO:10 or SEQ ID NO:12;
  - 5 (ii) comprises a fragment of SEQ ID NO:10 or SEQ ID NO:12 which fragment comprises at least a fragment of the extended portion of exon 2 of INSP100/INSP104 wherein the fragment of SEQ ID NO:10 or SEQ ID NO:12 functions as a growth hormone or has an antigenic determinant which is specific to the amino acid sequence as recited in SEQ ID NO.6; or
  - 10 (iii) is a functional equivalent of (i) or (ii).
2. A polypeptide according to claim 1 which:
  - (i) comprises the amino acid sequence as recited in SEQ ID NO:10 or SEQ ID NO:12;
  - (ii) comprises a fragment of SEQ ID NO:10 or SEQ ID NO:12 which fragment  
15 comprises at least a fragment of the extended portion of exon 2 of INSP100/INSP104 and wherein the fragment of SEQ ID NO:10 or SEQ ID NO:12 a growth hormone or has an antigenic determinant which is specific to the amino acid sequence as recited in SEQ ID NO.6; or
  - (iii) is a functional equivalent of (i) or (ii).
- 20 3. A polypeptide according to claim 1 or 2 which:
  - (i) consists of the amino acid sequence as recited in SEQ ID NO:10 or SEQ ID NO:12;
  - (ii) consists of a fragment of SEQ ID NO:10 or SEQ ID NO:12 which fragment  
25 comprises at least a fragment of the extended portion of exon 2 of INSP100/INSP104 and wherein the fragment of SEQ ID NO:10 or SEQ ID NO:12 functions as a growth hormone or has an antigenic determinant which is specific to the amino acid sequence as recited in SEQ ID NO.6; or
  - (iii) is a functional equivalent of (i) or (ii).
4. A polypeptide which is a functional equivalent according to part (iii) of any of the

above claims, characterised in that it is homologous to the amino acid sequence as recited in SEQ ID NO:10 or SEQ ID NO:12 and functions as a growth hormone or has an antigenic determinant which is specific to the amino acid sequence as recited in SEQ ID NO.6.

- 5 5. A polypeptide which is a fragment or a functional equivalent as recited in any one of claims 1 to 3, which has greater than 90% sequence identity with the amino acid sequence recited in SEQ ID NO:10 or SEQ ID NO:12, or with an active fragment thereof, preferably greater than 85%, 90%, 95%, 98% or 99% sequence identity.
- 10 6. A polypeptide which is a functional equivalent as recited in any one of claims 1 to 5, which exhibits significant structural homology with a polypeptide having the amino acid sequence recited in SEQ ID NO:10, or SEQ ID NO:12.
- 15 7. A polypeptide which is a fragment as recited in claims 1-3 and claim 5 having an antigenic determinant which is specific to the amino acid sequence as recited in SEQ ID NO.6 and which consists of 7, 9, 12 or more amino acid residues from the amino acid sequence recited in SEQ ID NO:10 or SEQ ID NO:12.
8. A purified nucleic acid molecule which encodes a polypeptide according to any one of the preceding claims.
9. A purified nucleic acid molecule according to claim 8, which comprises the nucleic acid sequence as recited in SEQ ID NO:9 or SEQ ID NO 11, or is a redundant  
20 equivalent or fragment thereof.
10. A purified nucleic acid molecule according to claim 8 which consists of the nucleic acid sequence as recited in SEQ ID NO:9 or SEQ ID NO:11, or is a redundant equivalent or fragment thereof.
11. A purified nucleic acid molecule which hybridizes under high stringency conditions  
25 with a nucleic acid molecule according to any one of claims 8 to 10.
12. A vector comprising a nucleic acid molecule as recited in any one of claims 8 to 11.
13. A host cell transformed with a vector according to claim 12.
14. A ligand which binds specifically to the growth hormone polypeptide according to any one of claims 1 to 7.
- 30 15. A ligand according to claim 14, which is an antibody.

16. A compound that either increases or decreases the level of expression or activity of a polypeptide according to any one of claims 1 to 7.
17. A compound according to claim 16 that binds to a polypeptide according to any one of claims 1 to 7 without inducing any of the biological effects of the polypeptide.
- 5 18. A compound according to claim 17, which is a natural or modified substrate, ligand, enzyme, receptor or structural or functional mimetic.
19. A polypeptide according to any one of claims 1 to 7, a nucleic acid molecule according to any one of claims 8 to 11, a vector according to claim 12, a host cell according to claim 13, a ligand according to claim 14 or claim 15, or a compound according to any  
10 one of claims 16 to 18, for use in therapy or diagnosis of disease.
20. A method of diagnosing a disease in a patient, comprising assessing the level of expression of a natural gene encoding a polypeptide according to any one of claims 1 to 7, or assessing the activity of a polypeptide according to any one of claims 1 to 7, in tissue from said patient and comparing said level of expression or activity to a control  
15 level, wherein a level that is different to said control level is indicative of disease.
21. A method according to claim 20 that is carried out *in vitro*.
22. A method according to claim 20 or claim 21, which comprises the steps of:
  - a) contacting a ligand according to claim 14 or claim 15 with a biological sample under conditions suitable for the formation of a ligand-polypeptide complex; and
  - 20 b) detecting said complex.
23. A method according to claim 20 or claim 21, comprising the steps of:
  - a) contacting a sample of tissue from the patient with a nucleic acid probe under stringent conditions that allow the formation of a hybrid complex between a nucleic acid molecule according to any one of claims 8 to 11 and the probe;
  - 25 b) contacting a control sample with said probe under the same conditions used in step a); and
  - c) detecting the presence of hybrid complexes in said samples; wherein detection of levels of the hybrid complex in the patient sample that differ from levels of the hybrid complex in the control sample is indicative of disease.

24. A method according to claim 20 or claim 21, comprising:
- a) contacting a sample of nucleic acid from tissue of the patient with a nucleic acid primer under stringent conditions that allow the formation of a hybrid complex between a nucleic acid molecule according to any one of claims 8 to 11 and the primer;
  - 5 b) contacting a control sample with said primer under the same conditions used in step a); and
  - c) amplifying the sampled nucleic acid; and
  - d) detecting the level of amplified nucleic acid from both patient and control samples; wherein detection of levels of the amplified nucleic acid in the patient sample that  
10 differ significantly from levels of the amplified nucleic acid in the control sample is indicative of disease.
25. A method according to claim 20 or claim 21 comprising:
- a) obtaining a tissue sample from a patient being tested for disease;
  - b) isolating a nucleic acid molecule according to any one of claims 8 to 11 from said  
15 tissue sample; and
  - c) diagnosing the patient for disease by detecting the presence of a mutation which is associated with disease in the nucleic acid molecule as an indication of the disease.
26. The method of claim 25, further comprising amplifying the nucleic acid molecule to form an amplified product and detecting the presence or absence of a mutation in the  
20 amplified product.
27. The method of claim 25 or claim 26, wherein the presence or absence of the mutation in the patient is detected by contacting said nucleic acid molecule with a nucleic acid probe that hybridises to said nucleic acid molecule under stringent conditions to form a hybrid double-stranded molecule, the hybrid double-stranded molecule having an  
25 unhybridised portion of the nucleic acid probe strand at any portion corresponding to a mutation associated with disease; and detecting the presence or absence of an unhybridised portion of the probe strand as an indication of the presence or absence of a disease-associated mutation.
28. A method according to any one of claims 20 to 28, wherein said disease includes, but is  
30 not limited to reproductive disorders, pregnancy disorder, such as gestational

- 5 trophoblastic disease, developmental disorders such as Silver-Russell syndrome, growth disorders, growth hormone deficiency, Cushing's disease, endocrine disorders, cell proliferative disorders, including neoplasm, carcinoma, pituitary tumour, ovary tumour, melanoma, lung, colorectal, breast, pancreas, head and neck, placental site
- 10 trophoblastic tumor, adenocarcinoma, choriocarcinoma, osteosarcoma and other solid tumours; angiogenesis, myeloproliferative disorders; autoimmune/inflammatory disorders; cardiovascular disorders; neurological disorders, pain; metabolic disorders including diabetes mellitus, osteoporosis, and obesity, cachexia, AIDS, renal disease; lung injury; ageing; infections including viral infection, bacterial infection, fungal
- 15 infection and parasitic infection, and other pathological conditions.
29. A method according to any one of claims 20 to 27, wherein said disease is a disease in which growth hormone proteins are implicated.
30. Use of a polypeptide according to any one of claims 1 to 7 as a growth hormone or as a modulator of growth hormone activity.
- 15 31. A pharmaceutical composition comprising a polypeptide according to any one of claims 1 to 7, a nucleic acid molecule according to any one of claims 8 to 11, a vector according to claim 12, a host cell according to claim 13, a ligand according to claim 14 or claim 15, or a compound according to any one of claims 16 to 18.
- 20 32. A vaccine composition comprising a polypeptide according to any one of claims 1 to 7 or a nucleic acid molecule according to any one of claims 8 to 11.
- 25 33. A polypeptide according to any one of claims 1 to 7, a nucleic acid molecule according to any one of claims 8 to 11, a vector according to claim 12, a host cell according to claim 13, a ligand according to claim 14 or claim 15, a compound according to any one of claims 16 to 18, or a pharmaceutical composition according to claim 31, for use in the manufacture of a medicament for the treatment of reproductive disorders, pregnancy disorder, such as gestational trophoblastic disease, developmental disorders such as Silver-Russell syndrome, growth disorders, growth hormone deficiency, Cushing's disease, endocrine disorders, cell proliferative disorders, including neoplasm, carcinoma, pituitary tumour, ovary tumour, melanoma, lung, colorectal, breast,
- 30 pancreas, head and neck, placental site trophoblastic tumor, adenocarcinoma, choriocarcinoma, osteosarcoma and other solid tumours; angiogenesis, myeloproliferative disorders; autoimmune/inflammatory disorders; cardiovascular

disorders; neurological disorders, pain; metabolic disorders including diabetes mellitus, osteoporosis, and obesity, cachexia, AIDS, renal disease; lung injury; ageing; infections including viral infection, bacterial infection, fungal infection and parasitic infection, and other pathological conditions.

- 5 34. A polypeptide according to any one of claims 1 to 7, a nucleic acid molecule according to any one of claims 8 to 11, a vector according to claim 12, a host cell according to claim 13, a ligand according to claim 14 or claim 15, a compound according to any one of claims 16 to 18, or a pharmaceutical composition according to claim 31, for use in the manufacture of a medicament for the treatment of a disease in which growth  
10 hormone proteins are implicated.
35. A method of treating a disease in a patient, comprising administering to the patient a polypeptide according to any one of claims 1 to 7, a nucleic acid molecule according to any one of claims 8 to 11, a vector according to claim 12, a host cell according to claim 13, a ligand according to claim 14 or claim 15, a compound according to any one of  
15 claims 16 to 18, or a pharmaceutical composition according to claim 31.
36. A method according to claim 35, wherein, for diseases in which the expression of the natural gene or the activity of the polypeptide is lower in a diseased patient when compared to the level of expression or activity in a healthy patient, the polypeptide, nucleic acid molecule, vector, ligand, compound or composition administered to the  
20 patient is an agonist.
37. A method according to claim 35, wherein, for diseases in which the expression of the natural gene or activity of the polypeptide is higher in a diseased patient when compared to the level of expression or activity in a healthy patient, the polypeptide, nucleic acid molecule, vector, ligand, compound or composition administered to the  
25 patient is an antagonist.
38. A method of monitoring the therapeutic treatment of disease in a patient, comprising monitoring over a period of time the level of expression or activity of a polypeptide according to any one of claims 1 to 7, or the level of expression of a nucleic acid molecule according to any one of claims 8 to 11 in tissue from said patient, wherein  
30 altering said level of expression or activity over the period of time towards a control level is indicative of regression of said disease.

39. A method for the identification of a compound that is effective in the treatment and/or diagnosis of disease, comprising contacting a polypeptide according to any one of claims 1 to 7 or a nucleic acid molecule according to any one of claims 8 to 11 with one or more compounds suspected of possessing binding affinity for said polypeptide or nucleic acid molecule, and selecting a compound that binds specifically to said nucleic acid molecule or polypeptide.
40. A kit useful for diagnosing disease comprising a first container containing a nucleic acid probe that hybridises under stringent conditions with a nucleic acid molecule according to any one of claims 8 to 11; a second container containing primers useful for amplifying said nucleic acid molecule; and instructions for using the probe and primers for facilitating the diagnosis of disease.
41. The kit of claim 40, further comprising a third container holding an agent for digesting unhybridised RNA.
42. A kit comprising an array of nucleic acid molecules, at least one of which is a nucleic acid molecule according to any one of claims 8 to 11.
43. A kit comprising one or more antibodies that bind to a polypeptide as recited in any one of claims 1 to 7; and a reagent useful for the detection of a binding reaction between said antibody and said polypeptide.
44. A transgenic or knockout non-human animal that has been transformed to express higher, lower or absent levels of a polypeptide according to any one of claims 1 to 7.
45. A method for screening for a compound effective to treat disease, by contacting a non-human transgenic animal according to claim 44 with a candidate compound and determining the effect of the compound on the disease of the animal.